

MODELING NCAA TOURNAMENT BRACKET POOL MULTIPLE ENTRY STRATEGY

Adams, Tom

www.poollogic.com

† E-mail: *tadamsmar@yahoo.com*

Single entry strategies for the NCAA tournament bracket pool have been explored in the statistical literature, but in-depth analysis has never been extended to multiple entry strategies. Modeling suggests the favorability of certain single entry strategies, but the estimated profits have never been directly confirmed in the most common real-world bracket pools: those that lack upset incentives. It is difficult to design a single-entry hypothesis test with high statistical power. Multiple entries represent one way to increase the statistical power.

A multiple entry strategy is developed and back-tested on 10 years of data from a real bracket pool. The strategy uses the following methods: a Markov model based on pre-tournament Sagarin ratings to estimate win probabilities, an opponent model based on contemporaneous opponent pick distribution data that is publicly available from large bracket contests (ESPN, Yahoo) before the tournament, Monte Carlo estimation of profit, hill-climbing optimization of profits. All these methods have been used in the literature to optimize brackets, but never in combination, and never to mutually optimize multiple brackets.

The profits or losses that would have accrued for 10 consecutive years (2008-2017) from entering 100 mutually optimized brackets in a real bracket pool with $\tilde{200}$ opposing brackets are measured and used to test the hypothesis that this strategy would have been profitable. The 10 year back-test yielded a 53% average yearly profit on exposed capital, with a one-tailed t -test p -value = 0.032.

IT'S A TOSS-UP: AN ANALYSIS OF COIN TOSS STRATEGY IN T20 CRICKET

Aradhyula, Sumhith Veda

Harvard University, Cambridge, MA, USA

† E-mail: *svaradhyula@college.harvard.edu*

As a quicker form of cricket than Test matches and One Day International (ODI) matches, the T20 format has emerged as a marketable form of the game in the 21st century. This format largely owes its popularity to the flashy, big money, and relatively recent Indian Premier League (IPL), considered the flagship tournament for T20 cricket. This study quantifies

the effects of (1) winning the coin toss and (2) the decision to bat or field made once a team wins the coin toss on the result of T20 games. This paper extends Dawson, et al.'s (Journal of the Operational Research Society, 2009) analysis of ODI matches to the very different T20 format. Data on 636 games from the first ten seasons (2008-2017) of the IPL were used in this analysis. Using linear probability and probit models, we analyzed home field advantage, pitch condition, and team quality metrics as determinants of the outcome of the game in addition to the coin toss. Results suggest that although the winner of the coin toss initially appears insignificant in determining the outcome of the given IPL game, at the 99% confidence level, the team that bats second (chasing team) has a significantly higher probability of winning. These results indicate that winning the coin toss could potentially prove a significant advantage, should captains improve their strategy in choosing whether to bat or chase.

BUMP, SET, SPIKE: USING ANALYTICS TO RATE VOLLEYBALL TEAMS AND PLAYERS

Bagley, Corley; Ware, Brett

VolleyMetrics, Provo, UT, USA

† E-mail: *corleybagley@gmail.com*

In the international arena, volleyball is one of the world's most popular sports. However, in U.S. collegiate volleyball, they don't get the resources as some of the other major sports, including football and basketball. As such, the methods used to analyze volleyball have been relegated to methods bordering on archaic. We have developed a system to rate teams and players based on their ability in various fundamentals within volleyball. Our initial goal was to provide a new way to grade passing other than the traditional three-or-four point scales that are currently used by most teams. This expanded to include similar analysis for the fundamentals of serving, attacking, blocking, and digging. By applying statistical methods to data provided by VolleyMetrics, we have developed an unbiased way to rate teams and individual players in each of these fundamentals. Our findings matched the actual results for men's collegiate volleyball last year. We also used these ratings to create a head-to-head model to predict the outcomes of matches with high accuracy. During two weeks of the 2016 men's season in which there were 52 matches played, our models predicted the winner with 87% accuracy. While these results are encouraging, we hope to improve our model with additional data that will allow us to be even more accurate in our ratings and, by result, our predictions.

USING SURVIVAL ANALYSIS TO ESTIMATE INJURY RECOVERY TIMES AND POSITIONAL VARIATION IN THE NFL

Binney, Zachary O (1); Janssens, A Cecile JW (1); Klein, Mitchel (1); Goodman, Michael (1); Hammond, Kyle E (2)

(1) *Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, USA*; (2) *Department of Orthopedics, Emory University School of Medicine, Atlanta, GA, USA*

† E-mail: *zbinney@emory.edu*

When an NFL athlete is injured the public relies on team-provided recovery time estimates. We sought instead to estimate recovery times for common NFL injuries, overall and by position, from historical injury report data. We defined time to injury recovery as the difference in weeks between an injury’s initial appearance on and its removal from weekly injury reports. We studied 5 common injuries from the 2007-2015 regular seasons using the Football Outsiders Injury Database. We built Kaplan-Meier (KM) curves for each position and injury category, with athletes censored at regular season’s end. The interquartile range (IQR) from each KM curve represents the time within which the middle half of injuries can be expected to resolve. In 17,310 player-seasons there were 2,334 ankle, 756 groin, 1,463 hamstring, 2,966 non-ACL knee, and 1,473 shoulder injuries. Median recovery was 2 weeks for all injuries. The proportion of injuries lasting ≤ 3 weeks ranged from 63% (knee, 95% confidence interval (CI) 61-65%) to 72% (groin, 95% CI 68-75%). IQR was 1-4 weeks for ankle, groin, and hamstring and 1-5 weeks for knee and shoulder injuries. Positional variation was greatest with shoulders: most positions had IQRs of 1-4 or 5 weeks, but defensive backs were 1-8 weeks, and there was insufficient follow-up to observe 75% of quarterbacks recovering. In summary, a majority of injuries resolved after 2 weeks, but recovery profiles varied by injury type and position. Limitations include heterogeneous injury categories and variable team injury reporting practices.

OPTIMIZING PRE-SEASON TRAINING LOADS IN AUSTRALIAN FOOTBALL

Carey, David L (1); Crow, Justin (1,2); Ong, Kok-Leong (1); Blanch, Peter (3); Morris, Meg E (1); Dascombe, Ben J (1); Crossley, Kay M (1)

(1) *La Trobe University, Melbourne, AUS*; (2) *Essendon Football Club, Melbourne, AUS*; (3) *Griffith University, Gold Coast, AUS*

† E-mail: *dlcarey88@gmail.com*

This study investigated whether pre-season training plans for Australian football can be computer generated using current training load guidelines to optimize injury risk reduction and performance improvement. A constrained optimization problem was defined for daily training loads in the pre-season phase of an elite Australian football team. Maximizing total

training volume and Banister model projected performance were considered as optimization objectives. Cumulative workload and acute:chronic workload ratio constraints were placed on training programs to reflect current guidelines on relative and absolute training loads for injury risk reduction. Optimization software was then used to generate pre-season training plans. The optimization framework was able to generate training plans that satisfied relative and absolute workload constraints. Increasing the off-season chronic training loads enabled prescription of higher amounts of ‘safe’ training and higher projected performance levels. Simulations showed that using a Banister model objective led to plans that included a taper in training load prior to competition in order to minimize fatigue and maximize projected performance. In contrast, when the objective was to maximize total training volume, more frequent training were prescribed in order to accumulate as much load as possible. An optimization approach provides an adaptable framework for physical preparation staff to quickly create athletic training plans that objectively optimize training goals whilst satisfying injury risk constraints, without exposing their plans to subjective bias. The method allows for individualized training plan design and the ability to adapt to changing training objectives and training load metrics.

USING A TENNIS RATING SYSTEM TO DETERMINE HANDICAPS IN AMATEUR MATCHES

Chan, Timothy CY (1); Singal, Raghav (2)

(1) *University of Toronto, Canada;* (2) *Columbia University, USA*

† E-mail: rs3566@columbia.edu

This paper builds on a recently developed Markov Decision Process-based (MDP) handicap system for tennis, which aims to make amateur/social matches more competitive. The system “gives” points to the weaker player based on skill difference, which is measured by the point-win probability. However, estimating point-win probabilities at the amateur level is challenging since point-by-point data is generally only available at the professional level. On the other hand, tennis rating systems are widely used and provide an estimate of the difference in ability between players, but a rigorous determination of handicap using rating systems is lacking. Therefore, our goal in this paper is to develop a mapping between the increasingly popular Universal Tennis Rating (UTR) system and the MDP-based handicaps, so that two amateur players can determine an appropriate handicap for their match based only on their UTRs. Using data from over 2500 ATP and WTA matches from 2015, we first develop and validate an approach to extract server-independent point-win probabilities from game-level data using the Markov chain model for a single game of tennis. Given the lack of point-level data for amateurs, we believe estimation of point-win probabilities from game-level data is important. Then, we show how to map server-independent point-win probabilities to server-specific point-win probabilities using Bayesian logistic regression. Finally, we use the estimated probabilities to produce handicaps via the MDP model, which

are regressed (Bayesian linear regression) against UTR differences between pairs of players. We conclude with thoughts on how a handicap system could be implemented in practice.

WHAT FACTORS IMPACTS A SUPERSTAR'S PERFORMANCE IN NBA?

Chen, Ruihao (1); Li, Mingfei (2)

(1) *Deloitte & Touche LLP, USA*; (2) *Bentley University, Waltham, MA, USA*

[†] E-mail: *anthonyruih.chen@gmail.com*

Nowadays, many sports superstars perform unbelievably well in their own fields. For example, NBA player Stephen Curry made 402 three pointers in the regular season 2015-16, obliterating the previous record of 286 three pointers in the regular season 2014-15. In addition, he had the second highest 3-point percentage in NBA with an accuracy of 44.4%. In this study, we explore the factors that may significantly impact the sports superstars' performances across multiple regular seasons. Variables include social media, team support, competitive opponents, etc. This study aims to support the superstar's team to mitigate the risk of a superstar's performance being fluctuated. Furthermore, opposing teams can leverage this study to predict a player's performance in upcoming games and strategize appropriately.

In this project, we take Stephen Curry as an example and analyze what impacts his 3-point performance, with regards to both quantity and accuracy. Data collected includes the overall performance of the Golden State Warriors (excluding Curry) per game, the opponent's overall performance of each game, Curry's performance per game, and Twitter data about him during the season 2015-16, etc. We applied Principal Component Analysis to identify key factors. Sentiment Analysis was also involved in order to understand the impact of social media on the player's performance. By implementing the Generalized Linear Model, we are able to identify significant factors on Curry's 3-point quantity and accuracy.

RAO-BLACKWELLIZING FIELD GOAL PERCENTAGE FOR NBA PLAYERS

Daly-Grafstein, Daniel; Bornn, Luke

Simon Fraser University, Burnaby, British Columbia, Canada

[†] E-mail: *danieldalygrafstein@gmail.com*

Shooting skill in the NBA is almost exclusively defined by field-goal percentage – the number of makes out of the total number of shots. Even advanced metrics like true shooting percentage still rely on counting the makes and misses of individual players. Here we attempt to better quantify a player’s shooting skill by using optical tracking data and shot trajectory characteristics to reduce the variability in predicting player field-goal percentages. Using tracking data, we model the trajectory of individual shots and use this to estimate shooting factors such as entry angle, shot depth, and left-right accuracy. We use these shooting factors to predict the probability of making a shot as well as measure shot accuracy bias and variation for individual players over the 2015-16 season. We then use these individual player measurements to stabilize inference of their field-goal percentage, allowing us to estimate metrics such as corner 3 shooting percentage earlier in a season than was previously possible. Finally, we present results on the effect of defense and minutes played on shot trajectory bias and variability, as well as differences in shooting factors for players over periods of hot and cold shooting.

A SURVEY OF ADVANCED MODELING TECHNIQUES FOR FORECASTING COLLEGE FOOTBALL GAME OUTCOMES

Egros, Edward (1); South, Charles (2)

(1) Fox 4 Sports, Dallas, TX; (2) UT Southwestern, Dallas, TX

† E-mail: *edward.egros@gmail.com*

There are many motives why data scientists and fans of college football would want to forecast the outcome of games – gambling, game prep (e.g. analyzing which factors a team should focus on), academic research, and so forth. As advanced statistical methods become more readily accessible, so do the opportunities to develop more robust forecasting models. Using data from the 2011 to 2015 seasons, we are implementing a variety of advanced modeling techniques to determine which best forecasts the outcome of games. These methods include ridge regression, the lasso, the elastic net, neural networks, random forests, k-nearest neighbors, gradient boosting machines and different Bayesian regression models. To evaluate the efficacy of the proposed models, we are testing them on data from the 2016 season; with all models accurately predicting the outcome at least 2/3 of the time, with the top performer – a Bayesian regression with team-specific variances – predicting the correct outcome 75.3% of the time.

PERFORMANCE VS. OBSERVATION: ARE EXPERTS AND ANALYSTS WATCHING THE SAME GAME?

Flaherty, Liam; Cleary, Richard

Babson College, Wellesley, MA, USA

† E-mail: *lflaherty1@babson.edu*

Baseball’s longstanding history of conflict between scouts and analysts came to light in popular culture with the successful 2003 release of Michael Lewis’ *Moneyball*. The argument presented in this project is that scouting feedback is not necessarily indicative of performance on the baseball field. To test this, we examined performance of pitchers from recent Babson College seasons, then assigned pairs of pitchers a “similarity score” of the type popularized by Bill James. This measure was based on the distance between the pair in terms of a standardized FIP-like measurement. We also designed a more qualitative approach in which three “scouts” (Babson Coaches) described the pitchers by choosing their outstanding attributes from a list of traditional baseball terms (i.e. command, control, velocity.) Pairs of pitchers were again ranked by similarity using a scoring system based on common attributes. In this initial study we did not find a strong association between performance similarity and scouting similarity, but the work does suggest ways we might find connections between scouts and analysts in professional data.

THE FIRST-PITCH GAME: STRATEGIES AND SIGNIFICANCE OF THE FIRST PITCH IN THE MLB

Gilfix, Adam; Morris, Carl; Avery, Christopher

Harvard University, Cambridge, MA, USA

† E-mail: *adam.gilfix@gmail.com*

There has been an immense amount of work achieved in better quantifying baseball players according to their contributions in almost every imaginable facet of the game. This has helped create a fairer market in the profession. However, there is less in-depth, publicly available research using newer, more detailed data, which lends itself to evaluating player strategies and the consequences thereof. Analysts have broken down the general importance of the first pitch and demonstrated the corresponding value of a first-pitch strike for a pitcher and first-pitch ball for batter. Similar research has revealed above-average results on first-pitch contact. And though other work has manifested strike- and swing-based tendencies that certain pitchers and hitters exhibit on 0-0 counts, there remains room to dig deeper into complex and motivating interaction-based analyses. Fluctuations in first-pitch strike and swing rates over the past few years have hinted at an equilibrium point, making this the perfect time to address the crucial 0-0 count. Previous studies are referenced and reanalyzed to give context to questions surrounding first-pitch significance and tactics. Hypotheses are formed and experimented on substantial data from TruMedia Networks, Inc., rejecting the notion of optimal mixed strategy equilibrium in 0-0 counts and demonstrating the opportunity for differing areas of strategical exploitation and improvement on the first pitch. We find that pitchers should be throwing more first-pitch strikes and batters should swing more at first pitches over the heart of the plate.

COMPARING GOLF ERAS: AN ANALYSIS OF GOLF'S BEST TO DETERMINE THE GREATEST PROFESSIONAL GOLFER OF ALL TIME

Gravelle, Patrick

Queen's University, Kingston, Ontario, Canada

† E-mail: *patrick.gravelle@queensu.ca*

Competition can be found in a schoolyard baseball game, a retirement home shuffleboard match, an NBA championship final, and anywhere in between. It is through this competition that the motivation to best one's opponent is found, and underlying that motivation, is the desire to be the greatest. But in sports, this theme of greatness does not simply extend to just one match, or a single season, but rather, it reaches throughout all of time. For a sport such as golf, like many others, the debate of determining the greatest player of all time appears hopelessly divided. This paper aims to close that division by analyzing several of the major statistical categories for professional golfers, in which total event wins and win percentage is ranked for both PGA and Major Tournaments. In addition, two new categories will be evaluated, comparing the elite golfers amongst one another by equating their respective results for every Major Tournament event participated in as a professional. This allows for a direct comparison between golfers of different eras as it will be assumed that the first time playing in an event is of equal difficulty for each player, regardless of the year. Each distinct categorical assessment is given a weighted value which corresponds to the respective category's level of importance. These evaluations, combined with their weighted values, produce an overall score in which the golfers are then ranked in descending order to yield the final results.

PROJECTING COLLEGE BASKETBALL FRESHMAN PERFORMANCE USING GRASSROOTS BASKETBALL STATISTICS

Johnson, Neil

ESPN, Bristol, CT, USA

† E-mail: *neil.johnson@espn.com*

Basketball recruiting relies largely, if not solely, on the opinions of scouts who travel the country to see and evaluate as many players as they can. Logistically, it is impossible for scouts to see every player that will play National Collegiate Athletic Association (NCAA)

Division 1 basketball, and there is a bias towards evaluating the best players and teams with more exposure. With the rise of Grassroots leagues sponsored by athletic apparel companies, prospective basketball recruits now play in organized leagues that level the playing field and provide consistent statistics. Using available box scores statistics, this paper will create Box Plus/Minus ratings for the players in those Grassroots leagues and perform polynomial regression between the Grassroots and College Box Plus/Minus ratings for all players who participated at both levels. This paper will then summarize the results by evaluating the utility of forecasting the performance of NCAA Division 1 freshman basketball players given their Grassroots Box/Plus Minus ratings.

NEW WAYS TO EVALUATE PITCHERS USING WEIGHTED OFFENSIVE AVERAGE AND PENALIZING EARNED RUNS

Kim, Chulmin

University of West Georgia, Carrollton, GA, USA

[†] E-mail: *ckim@westga.edu*

For measuring a pitcher's performance Earned run average (ERA) has been a dominant statistic to along with number of wins (W), walks plus hits per inning pitched (WHIP), and opponents' batting average (OBA). ERA is the mean of earned runs given up by a pitcher per nine innings pitched. Weighted offensive average (WOA), explains a batter's overall performance to generate runs and introduced by Kim (2013), is used to develop the opponents' batting average (oWOA) as an alternative of ERA.

ERA roles differently between starting and relief pitchers. If a relief pitcher enters the game with scoring position and allows a hit which scores runs, he is not charged with those runs. Also relief pitchers typically pitch for short innings so they can more concentrate on each pitch, unlike starters who need to save their energy over 6 or 7 innings. This may lead the relievers keep their ERAs down. We modify the current ERA by penalizing relief pitchers' earned runs even they enter the games already with scoring positions. We propose allocating earned runs based on where runners were left on base when a pitcher is relieved. We allocate 1/4, 2/4, and 3/4 of an earned run to the starter for runners left on the 1st, 2nd, and 3rd base, respectively, that score while the reliever is pitching. The remaining of the earned run goes to the reliever. We evaluate the top five 'oWOA' MLB pitchers in 2016 using this new ERA.

EXPERIENCED VETERAN OR PROMISING ROOKIE: THE EFFECTS OF MLB MANAGERS' SALARY AND EXPERIENCE ON TEAM SUCCESS

Kim, J.C. (1); Kim, Seong Dae (2)

(1) *Fairleigh Dickinson University, Teaneck, NJ, USA*; (2) *University of Alaska Anchorage, Anchorage, AK, USA*

† E-mail: *jckim1@fdu.edu*

Based on a 2007 article in Sport Business Daily, MLB managers averaged a \$1.4 M salary in 2007. For the 2017 season, an average salary of the top-five highest-paid managers is a \$4.5 M. Unsurprisingly, four out of five managers are employed with teams located in large cities such as New York, L.A., Chicago, and San Francisco. Also, the five managers have many years of experience as a manager. It is no doubt that a manager’s job responsibilities and duties are quite significant and their salary is considered an expense from the team’s perspective. Thus, hiring a new manager could be a more difficult decision to teams in small markets than to those in big markets. Teams in small markets may want to consider hiring a more affordable and less experienced but promising young manager to save money to hire good players, rather than an expensive and experienced one. Given the MLB managers’ salaries and its business challenge, we offer the following research questions: 1) Do teams with experienced managers perform better? 2) Do teams with high-paid managers perform better? To answer the questions above, we will examine the effects of MLB managers’ salaries and their experiences on their teams’ success. We will come up with different ways to measure the team’s success and explore manager-related factors that might be positively or negatively related to the team’s performance.

A TIMEOUT IS A TIMEOUT? EFFECTS OF A RULE CHANGE IN THE NATIONAL HOCKEY LEAGUE (NHL)

Davis, Jack (1), Kniffin, Kevin M (2), Hilbrich, Christian (2)

(1) *Simon Fraser University, Burnaby, British Columbia, Canada*; (2) *Cornell University, Ithaca, NY, USA*

† E-mail: *kmk276@cornell.edu*

Starting with the 2015-16 National Hockey League (NHL) season, coaches have been able to risk their one timeout allocated per game to challenge goals if they contended the other team either interfered with the goalie or were offside en route to the goal. In order to examine the impact of this rule change, we analyzed play-by-play records for the two seasons before and the two seasons after the rule change. We find that when timeouts can be used by coaches to challenge a scoring play, there are (1) significantly fewer timeouts called and (2) those timeouts that are used tend to be called later in games. In addition, we find that (3) the odds of a timeout being called immediately after an opponent scores have approximately doubled since the adoption of the rule change, presumably because it is the main window of time when a timeout can now be put on the line to challenge a goal. Our study contributes to a better understanding of the effects of rule changes in sports as well as why coaches use timeouts.

GRIP IT AND RIP IT: A FURTHER EXPLORATION OF PREDICTING SUCCESS FOR MALE PROFESSIONAL GOLFERS ON THE PGA TOUR

Machiorlatti, Michael; Dvorak, Justin

University of Oklahoma Health Sciences Center

† E-mail: *michael-machiorlatti@ouhsc.edu*

Success in golf is hypothesized to be influenced by the ability to be a long ball hitter, having command of one's irons, and having a feathery touch around the greens. Prior studies have explored these common beliefs in order to predict golf success. This study continued with the trend of most previous work (Berry 1999, Nero 2001, Rishe 2001, Alexander and Kern 2005, Kahan 2010, Baugher 2016), which used earnings as the metric to measure golf success; however, we also explored salary rank by year. Data were obtained from ESPN, CBS Sports, and the PGA Tour, and incorporated traditional shot measures in the modeling process. This study expanded upon previous studies by considering a wider range of covariate transformations, using more specific shot measures, exploring data over a greater time frame (2003 to 2016), and incorporating longitudinal analysis into the modeling process. While our results were consistent with prior studies, finding scoring average to be a major contributor to success, we were also able to find more specific predictors of golfers' earnings over time.

FROM NOOB TO SMURF: ADVANCED ANALYTICS FOR LEAGUE OF LEGENDS

Maymin, Philip Z.

University of Bridgeport, Bridgeport, CT, USA and Vantage Sports

† E-mail: *philip@maymin.com*

Standard metrics for multiplayer online battle arena (MOBA) games like League of Legends (LoL) are very simple: kills, deaths, and the like. At Vantage Sports, we use a proprietary method to generate unique metrics that are more useful for professional players. These metrics are then calculated for hundreds of thousands of amateur player games, and the results used to determine which ones most contribute to winning. Some of the most important ones are worthless deaths and smart kills, which refine the standard metrics based on whether the team overall benefited from the activity. A new player rating model described here correlates strongly with winning even though it is essentially based on just one individual's contribution to a five-on-five game.

FROM MARKOV MODELS TO POISSON POINT PROCESSES: UNDERSTANDING PLAYER MOVEMENT IN THE NBA

Mortensen, Jacob; Bornn, Luke

Simon Fraser University, Burnaby, British Columbia, Canada

† E-mail: jmortens@sfu.ca

When considering movement in space, a useful tool is a Markov model, where the position of the agent at time $t + 1$ depends only on their position at time t . In this paper we build on existing theory to show that as the number of spatial locations in a bounded region approaches infinity, a Markov model can be represented by a Poisson point process, a popular type of spatial model that accounts for correlation between nearby locations. Using SportVu player tracking data provided by the National Basketball Association we show how this relationship can be leveraged to produce distinct maps of player movement for each team in the NBA. By comparing these maps, we can understand the tendencies of each team and how teams utilize court space differently.

ISLANDS IN THE SEA ARE CONNECTED IN THE DEEP: PROFESSIONAL SOCCER PLAYER TRANSFERS IN BIG 5 EUROPEAN LEAGUES USING EXPONENTIAL RANDOM GRAPH MODEL (ERGM)

Mukherjee, Subhasree (1); Dhayanithy, Deepak (2)

(1) *Indian Institute of Management Kozhikode, Kerala, India* (2) *Indian Institute of Management Kozhikode, Kerala, India*

† E-mail: subhasreem08fpm@iimk.ac.in

The 2016 revenue of the Big 5 European soccer leagues (England, France, Spain, Italy, and Germany) was £ 12.89 Billion (compared to NFL's annual revenue of £ 12.27 Billion). Combined valuation of players in just one league (England) was £ 4.98 Billion. Player transfers amongst these clubs are strongly inter-related, thus making transfer market a clear case of violation of the 'independent observations' assumption of conventional regression models. For instance, Pogba's move from Juventus to Manchester United in 2015 is clearly related to Higuain's move from Napoli to Juventus amongst other player purchases of Juventus. Extant literature of player transfers does not address this serious issue. Grounded in Strategic Management literature, we use Resource Dependence Theory to investigate tie-formation

in player transfer network involving European Big 5 Soccer Leagues, using ERGM. ERGM captures the dynamic evolution of a network in combination with nodal (club) attributes and network structural elements. 2015 Player transfer data for this study is obtained from transfermarkt.com, fifaindex.com, and Deloitte Football Money League. Clubs' previous season standings, network status, resource richness (financial), squad rating (FIFA player rating), and squad size are significant explanatory variables under the 'independent observations' assumption. However, after incorporating network dynamics and interrelationships in ERGM, only previous season standing remain to be significant. Network properties such as edges, cyclical ties and popularity are significant determinants of professional soccer player transfers. Network perspective using ERGM is a more realistic framework to understand drivers of soccer player transfers as compared to conventional regression models.

WHAT DOES IT MEAN TO DRAFT PERFECTLY?: A GAME THEORETIC FRAMEWORK FOR THE NHL ENTRY DRAFT

Nandakumar, Namita

University of Pennsylvania, Philadelphia, PA, USA

[†] E-mail: *namitan@wharton.upenn.edu*

This project defines what it means for a team to be successful in drafting and determines if any NHL teams exhibited a sustained competitive edge in drafting from 2000 to 2009. At a high level, we compare actual draft outcomes to teams' perfect draft outcomes. A perfect draft can be thought of as what would happen if a GM could redo a draft with complete knowledge of prospects' career values as well as other teams' choices, which remain unchanged. In this scenario, it is optimal to select good players just before other teams do.

We use this as a benchmark for a draft efficiency metric, which retrospectively analyzes how well teams drafted relative to their best case scenario in each year. This allows us to compare draft outcomes fairly, and we find that no NHL team had a statistically significant advantage in extracting value from the draft over the ten-year period. However, teams like the Blackhawks set themselves up for success with many high-value picks.

Teams can prospectively utilize this framework by modeling two inputs for every eligible player: expected NHL value as well as a probability distribution of possible draft positions. They can then algorithmically define and order a series of selections that maximizes total expected NHL value. The advantage of this approach over simply drafting the best player available at each pick fades in later rounds. Still, there are notable recent instances where teams would have theoretically benefited from waiting longer to pick their preferred prospects.

SCHEDULING EFFECTS IN THE NBA AND NHL AND THEIR BETTING MARKETS

Osborne, Jason A Osborne; Post, Justin B.

North Carolina State University, Raleigh, NC, USA

[†] E-mail: *jaosborn@ncsu.edu*

Teams in the NBA have are giving increasing consideration to the issue of fatigue, recognizing that without rest, performance might be compromised. Using generalized linear mixed models we estimate effects of schedule conditions, whether home/road teams were idle, home or away the day prior to a regular season contest in professional hockey and basketball. We attempt to control for team strengths with factorial effects for home and visiting teams. On the scale of win probabilities, we find the magnitude of the estimated effects of one day of rest for the home team and the away team to be highly significant, additive and comparable across the two sports. The relative change in estimated win probability from the least to the most favorable schedule conditions is as large as 17% (NHL) or 18% (NBA).

We also find evidence that bettors are taking schedules into consideration. Published point spreads and over/under (o/u) lines in the NBA differ significantly across schedule conditions, achieving an ordering that is consistent with the conventional wisdom that having to play the day before a contest and then travel has a negative effect on performance. Interestingly, the observed game totals do not differ significantly across scheduling conditions, nor reflect this conventional wisdom. Lastly, inclusion of an overtime effect can “debias” factorial team effect estimates and improve predictions made prospectively (without knowledge of whether wagered-upon games went into overtime), identifying a potential market inefficiency.

RISK FACTORS AND PREDICTIVE MODELING OF THE UCL INJURY USING PITCHF/X DATA IN MLB

Sakaori, Fumitake (1); Hoshina, Ibuki (2)

(1) Chuo University, Tokyo, Japan; (2) Shiga University, Shiga, Japan

[†] E-mail: *sakaori@math.chuo-u.ac.jp*

Ulnar collateral ligament (UCL) reconstruction surgeries, also known as Tommy John surgeries (TJS), for baseball pitchers have increased in recent decades. The UCL injury can be caused by in the iterative stress to the UCL in the medial elbow. In spite of the importance of preventing UCL injuries, the scientific consensus of the risk factors for the injuries has never been obtained yet. In this presentation, we reconsider candidates of risk factors for the injuries referring to some experts’ opinions, and then obtain adjusted odds ratios for selected risk factors by a logistic regression model. As a result, for starter pitchers, a smaller repertoire of pitch types, a farther horizontal release location away from the body, and a greater mean pitch count per game are selected risk factors. Also, for relief pitchers, a smaller

repertoire of pitch types, a farther horizontal release location away from the body, a greater mean pitch speed of fast balls, and fewer days between consecutive games are selected. Also, we consider some predictive modeling methods to predict the UCL injuries.

AN ANALYSIS OF SHOT QUALITY IN MAJOR LEAGUE LACROSSE (MLL)

McNulty, Jesse (1); Long, Christopher (2); Schanzenbach, Tyler G (3); Keegan, Joseph (4)

(1) Atlanta Blaze, Kennesaw, GA, USA; (2) Vorpala, LLC., Versailles, KY, USA; (3) William Penn High School, New Castle, DE, USA; (4) U.S. Lacrosse Magazine, Boston, MA, USA

† E-mail: jessemcnult@gmail.com

Analytics in the game of professional field lacrosse is very much in its infancy. This paper analyzes spatial and visual tendencies as means to enhance lacrosse expertise. We introduce analytical techniques designed to quantify, visualize, and communicate spatial aspects of Major League Lacrosse (MLL) performance. We propose mixed effects modeling to quantify shooting quality of MLL players and present practices that differentiate MLL players' shooting ability. We conduct a case study to inspect shot site performances for every player in Major League Lacrosse and determine which players exhibit the most potent offensive behaviors. We conclude by proposing that visual and spatial analysis represent new methodologies for analysis of professional field lacrosse.

PREDICTING SHOT SUCCESS USING REAL ESTATE

Seward, Neil (1); Loyzer, Mark (2); Makrehchi, Masoud (1)

(1) University of Ontario Institute of Technology, Oshawa, Ontario, Canada (2) University of Toronto, Toronto, Ontario, Canada

† E-mail: neil.seward@uoit.ca

Neural networks have been used in myriad prediction tasks and outperform other non neural network based models. One area where learning tasks, in general, have not been extensively used is in adversarial multi-agent trajectory domains. Traditionally, there has not been a natural way to represent trajectory inputs which are continuous and spatially correlated that are amenable to existing machine learning techniques. For the problem of predicting whether a basketball shot is a make or a miss in the National Basketball Association (NBA), different methods have been used in representing players trajectories for prediction tasks. We

investigate exploiting the temporal information in trajectories for purposes of shot prediction. Further exploring trajectory representations and player movement, we leverage additional information to represent as spatial features within polygons that represent space that is owned by a player. For offensive spatial features present in the polygons, channel shading can represent shooting ability over the space that is owned by the shooter. We also present a framework to identify and quantify defensive congestion during a shot to create value behind defensive positioning. Finally we assign player roles through channels formed from encoded player representations. By representing different spatial features of player movement during a possession, we aim to provide increased accuracy when determining if a basketball shot is a make or a miss.

QUANTIFYING THE CAUSAL EFFECTS OF CONSERVATIVE FOURTH DOWN DECISION MAKING IN THE NATIONAL FOOTBALL LEAGUE

Yam, Derrick (1); Lopez, Michael (2)

(1) *Brown University, Providence, RI, USA*; (2) *Skidmore College, Saratoga Springs, NY, USA*

[†] E-mail: *yam.derrick@gmail.com*

It is often argued that football coaches are too conservative in attempting fourth down conversions. However, coaches' decisions are not randomized, and because teams going for it do so by choice, traditional research into team decision-making may be confounded by extraneous factors. For example, teams going for it generally do so because they are trailing, and thus inference on how all teams should behave has required unjustifiable extrapolations. Using the National Football League (NFL), we attempt to estimate the causal effects of implementing a more aggressive fourth down strategy by approximating the additional number of wins that any one NFL team could have gained. Utilizing a nearest neighbor matching algorithm, teams that went for it are paired to those who did not go for it based on their probability of going for it, defined as the propensity score, as well as other game-level factors. By looking at outcomes within the cohort of matched plays, we find that, on average, imposing a simple, more aggressive fourth down strategy would have added an estimated 3.3 wins across 13 seasons. Our results better inform decision-making in a high-stakes environment where standard statistical tools are informative but, to date, limited.